



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b>  <b>A61K</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/18353</b>  <b>(43) International Publication Date:</b> 6 April 2000 (06.04.00)
<b>(21) International Application Number:</b> PCT/US99/22354 <b>(22) International Filing Date:</b> 28 September 1999 (28.09.99)  <b>(30) Priority Data:</b> 60/102,056 28 September 1998 (28.09.98) US  <b>(71) Applicant (for all designated States except US):</b> THE RE- SEARCH FOUNDATION OF STATE UNIVERSITY OF NEW YORK [US/US]; P.O. Box 9, Albany, NY 12201-0009 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> RYAN, Maria, Emanuel [US/US]; 5 Vannina Place, Huntington, NY 11743 (US). GOLUB, Lorne, M. [US/US]; 29 Whitney Gate, Smith- town, NY 11787 (US). RAMAMURTHY, Nungavaram, S. [US/US]; 10 Lynam Court, Smithtown, NY 11787 (US).  <b>(74) Agent:</b> BARON, Ronald, J.; Hoffmann & Baron, LLP, 6900 Jericho Turnpike, Syosset, NY 11791 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished          upon receipt of that report.</i>
<b>(54) Title:</b> A NOVEL INHIBITOR OF CATARACT FORMATION  <b>(57) Abstract</b>  Methods of reducing the risk of cataract development in a mammal are provided and include administering to the mammal an effective amount of a tetracycline derivative. A preferred tetracycline derivative administered according to the methods of the present invention is 6 $\alpha$ -deoxy 5-hydroxy-4-dedimethylaminotetracycline.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	RS	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## **A NOVEL INHIBITOR OF CATARACT FORMATION**

This invention was made with support from the United States government. The United States government has certain rights in this invention.

5

### **BACKGROUND OF THE INVENTION**

Cataracts are an abnormal progressive condition of the lens of the eye, characterized by loss of transparency. A gray-white opacity can be seen within the lens, behind the pupil. Most cataracts are caused by degenerative changes, occurring most often after fifty years of age, however in susceptible patients such as diabetics, cataracts can occur at a much earlier age.

10

Two types of cataracts have been described: (1) metabolic or juvenile cataracts, which are observed in children and young adults with uncontrolled diabetes, and (2) senile cataracts which are more common than metabolic cataracts. The cataracts in diabetics are similar to the senile cataracts observed in non-diabetic patients but tend to occur at a younger age (e.g. the aging process is accelerated).

15

The tendency to develop cataracts is inherited and can be accelerated by diseases such as diabetes. If cataracts are untreated, sight is eventually lost. First, vision is blurred, then bright lights glare diffusely, and finally, distortion and double vision may develop.

20


Previously, cataracts have been treated according to the progressive condition of the lens of the eye. In particular, senile cataracts are usually treated with excision of the lens and prescription of special contact lenses or glasses. The soft cataracts of children and young adults can either be incised and drained or fragmented by ultrasound, followed by irrigation and aspiration of the fragments through a minute incision.

25


As stated above, the tendency to develop cataracts is inherited and can be accelerated by diseases such as diabetes. Diabetes is the leading cause of blindness in adults and accounts for loss of vision in 8% of those who are legally blind in the United States (Klein and Klein, 1995). Sixty-five percent of diabetic patients develop blindness within five years after detection of proliferative retinopathy (Lavine, 1990). An additional complication of diabetes in the eye is associated with swelling of the lens. This swelling is a result of the accumulation of fructose and sorbitol that increases the osmolality within the lens. As this process continues, lens protein becomes denatured and cataracts form (Lavine, 1990).

Studies of the blood vessels in the eye of an animal model of Type II diabetes, the ZDF/Gmi-*fa* rat, have demonstrated microscopic vascular changes in the retina, such as an increase in capillary nuclear density and an increase in basement membrane thickness, when compared to non-diabetic control animals (Danis *et al.*, 1993). Danis *et al.*, demonstrated that these animals develop microvascular retinopathy determined morphologically by increases in the numbers of endothelial cells (endothelial hyperplasia) and associated capillaries. A prominent early marker for diabetic retinopathy is pericyte degeneration and the development of pericyte ghosts. Another eye complication seen in this diabetic rat model is the development of cataracts.

Using compounds approved for use in humans and available in the oral, injectable and topical routes, such as tetracycline, to treat diseases or conditions such as cataracts has not been suggested. The compound, tetracycline, exhibits the following general structure:



The numbering system of the ring nucleus is as follows:



Tetracycline as well as the 5-OH (Terramycin) and 7-Cl (Aureomycin) derivatives exist in nature, and are well known antibiotics. Natural tetracyclines can be modified without losing their antibiotic properties, although certain elements of the structure must be retained. The modifications that can and cannot be made to the basic tetracycline structure have been reviewed by Mitscher in *The Chemistry of Tetracyclines*, Chapter 6, Marcel Dekker, Publishers, New York (1978). According to Mitscher, the substituents at positions 5-9 of the tetracycline ring system may be modified without the complete loss of antibiotic properties. Changes to the basic ring system or replacement of the substituents at positions 1-4 and 10-12, however, generally lead to synthetic tetracyclines with substantially less or effectively no antimicrobial activity. An example of a chemically modified tetracycline (hereinafter CMT) is 4-dedimethylaminotetracycline which is commonly considered to be a non-antimicrobial tetracycline.

In addition to their antibiotic properties, tetracyclines have been described as having a number of other uses. For example, tetracyclines are also known to inhibit the activity of collagen destructive enzymes such as mammalian collagenase, gelatinase, macrophage elastase and bacterial collagenase. Golub et al., *J. Periodont. Res.* 20:12-23 (1985); Golub et al. *Crit. Revs. Oral Biol. Med.* 2: 297-322 (1991); U.S. Pat. Nos. 4,666,897; 4,704,383; 4,935,411; 4,935,412. In addition, tetracyclines have been known to inhibit wasting and protein degradation in mammalian skeletal muscle, U.S. Pat. No. 5,045,538.

Furthermore, tetracyclines have been shown to enhance bone protein synthesis in U.S. Pat. No. Re. 34,656 and to reduce bone resorption in organ culture in U.S. Pat. No. 4,704,383.

5 Similarly, U.S. Pat. No. 5,532,227 to Golub et al., discloses that tetracyclines can ameliorate the excessive glycosylation of proteins. In particular, tetracyclines inhibit the excessive collagen cross linking which results from excessive glycosylation of collagen in diabetes.

10 These properties cause tetracyclines to be useful in treating a number of diseases. For example, there have been a number of suggestions that tetracyclines, including non-antimicrobial tetracyclines, are effective in treating arthritis. See, for example, Greenwald et al., "Tetracyclines Suppress Metalloproteinase Activity in Adjuvant Arthritis and, in Combination with Flurbiprofen, Ameliorate Bone Damage," *Journal of Rheumatology* 19:927-938(1992); Greenwald et al., "Treatment of Destructive Arthritic Disorders with MMP Inhibitors: Potential Role of

15 Tetracyclines in, Inhibition of Matrix Metalloproteinases: Therapeutic Potential," *Annals of the New York Academy of Sciences* 732: 181-198 (1994); Kloppenburg et al., "Minocycline in Active Rheumatoid Arthritis," *Arthritis Rheum* 37:629-636(1994); Ryan et al., "Potential of Tetracycline to Modify Cartilage Breakdown in Osteoarthritis," *Current Opinion in Rheumatology* 8: 238-247(1996); O'Dell et al.,

20 "Treatment of Early Rheumatoid Arthritis with Minocycline or Placebo," *Arthritis Rheum* 40:842-848(1997).

Tetracyclines have also been suggested for use in treating skin diseases. For example, White et al., *Lancet*, Apr. 29, p. 966 (1989) reports that the tetracycline minocycline is effective in treating dystrophic epidermolysis bullosa, which is a life-

25 threatening skin condition believed to be related to excess collagenase.

The effectiveness of tetracycline in skin disorders has also been studied by Elewski et al., *Journal of the American Academy of Dermatology* 8:807-812 (1983). Elewski et al. disclosed that tetracycline antibiotics may have anti-inflammatory activity in skin diseases.

5            Similarly, Plewig et al., *Journal of Investigative Dermatology* 65:532 (1975), disclose experiments designed to test the hypothesis that antimicrobials are effective in treating inflammatory dermatoses. The experiments of Plewig et al. establish that tetracyclines have anti-inflammatory properties in treating pustules induced by potassium iodide patches.

10           The use of tetracyclines in combination with non-steroidal anti-inflammatory agents has been studied in the treatment of inflammatory skin disorders caused by acne vulgaris. Wong et al., *Journal of American Academy of Dermatology* 1: 1076-1081 (1984), studied the combination of tetracycline and ibuprofen and found tetracycline to be an effective agent against acne vulgaris and ibuprofen to be useful in  
15           reducing the resulting inflammation by inhibition of cyclooxygenase. Funt et al., *Journal of the American Academy of Dermatology* 13: 524-525 (1985), disclosed similar results by combining antimicrobial doses of minocycline with ibuprofen.

             An antimicrobial tetracycline derivative, doxycycline, has been used to inhibit nitrate production. D'Agostino et al., *Journal of Infectious Diseases*: 177:489-92  
20           (1998), discloses experiments where doxycycline, administered to mice injected with bacterial lipopolysaccharide (hereinafter LPS), exerted a protective effect by inhibiting nitrate production by an IL-10 independent mechanism. Experiments carried out *in vitro* also showed that doxycycline inhibited nitric oxide synthesis by LPS activated macrophages without enhancing endogenous IL-10 release.

25

             Based on the foregoing, tetracyclines have been found to be effective in different treatments. However, there has been no suggestion whatsoever that tetracyclines can be used to reduce the risk of cataract development in a mammal.

Accordingly, it is one of the purposes of this invention, among others, to provide an economical and relatively uncomplicated method of reducing the risk of cataract development.

### SUMMARY OF THE INVENTION

5 It has now been discovered that these and other purposes can be achieved by the present invention, which provides for methods of reducing the risk of cataract development in a mammal.

The methods of the present invention provide for reducing the risk of cataract development in a mammal by administering to the mammal an effective amount of a  
10 tetracycline derivative. A preferred method of the present invention provides that the tetracycline derivative is a non-antimicrobial tetracycline.

Another preferred method of the present invention provides for reducing the risk of cataract development by administering to a mammal an effective amount of a chemically-modified tetracycline derivative, for example  
15 dedimethylaminotetracycline. A preferred dedimethylaminotetracycline derivative is 6 $\alpha$ -deoxy 5-hydroxy 4-dedimethylaminotetracycline (CMT-8).

Other dedimethylaminotetracyclines which can be administered according to the methods of the present invention include 4-dedimethylaminotetracycline, 4-dedimethylamino-5-oxytetracycline, 4-dedimethylamino-7-chlorotetracycline, 4-  
20 hydroxy-4-dedimethylaminotetracycline, 5a, 6-anhydro-4-hydroxy-4-dedimethylaminotetracycline, 6  $\alpha$  -deoxy-5-hydroxy-4-dedimethylaminotetracycline, 6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 4-dedimethylamino-12a-deoxytetracycline, 12 $\alpha$  -deoxy-4-deoxy-4-dedimethylaminotetracycline, 12a, 4  $\alpha$  -  
25 anhydro-4-dedimethylaminotetracycline, 7-dimethylamino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 5-hydroxy-6-  $\alpha$  -deoxy-4-dedimethylaminotetracycline, 4-dedimethylamino-12  $\alpha$  -deoxyanhydrotetracycline and 4-dedimethylamino-11-hydroxy-12a-deoxytetracycline.

Other dedimethylaminotetracyclines which can be administered according to the methods of the present invention have D ring substituents at the C7 and/or C9 positions on the 4-dedimethylaminotetracycline molecule. These compounds include

7-azido-6-demethyl-6-deoxy-4-dedimethylamino tetracycline, 7-dimethylamino-9-azido-6-demethyl-6-deoxy-4-dedimethylamino tetracycline, 9-amino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 9-azido-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 9-nitro-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 9-amino-7-dimethylamino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 7-acetamido-6-demethyl-6-deoxy-4-dedimethylamino tetracycline, 9-acetamido-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 7-amino-9-nitro-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 9-(N,N-dimethyl)glycylamino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 7, 9-diamino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 9-ethoxythiocarbonylthio-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 7-dimethylamino-9-acetamido-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 9-azido-7-dimethylamino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 9-azido-7-dimethylamino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 7-azido-5-hydroxy-6-deoxy-4-dedimethylamino tetracycline, 7-dimethylamino-9-azido-5-hydroxy-6-deoxy-4-dedimethylamino tetracycline, 9-amino-5-hydroxy-6-deoxy-4-dedimethylaminotetracycline, 9-azido-5-hydroxy-6-deoxy-4-dedimethylaminotetracycline, 9-nitro-5-hydroxy-6-deoxy-4-dedimethylaminotetracycline, 9-amino-7-dimethylamino-5-hydroxy-6-deoxy-4-dedimethylaminotetracycline, 7-acetamido-5-hydroxy-6-deoxy-4-dedimethylamino tetracycline, 9-acetamido-5-hydroxy-6-deoxy-4-dedimethylaminotetracycline, 7-amino-9-nitro-5-hydroxy-6-deoxy-4-dedimethylaminotetracycline, 9-(N,N-dimethyl)glycylamino-5-hydroxy-6-deoxy-4-dedimethylaminotetracycline, 7, 9-diamino-5-hydroxy-6-deoxy-4-dedimethylaminotetracycline, 7-dimethylamino-9-amino-5-hydroxy-6-deoxy-4-dedimethylaminotetracycline, 9-ethoxythiocarbonylthio-5-hydroxy-6-deoxy-4-dedimethylaminotetracycline, 7-dimethylamino-9-acetamido-5-hydroxy-6-deoxy-4-dedimethylaminotetracycline, 9-azido-7-dimethylamino-5-hydroxy-6-deoxy-4-dedimethylaminotetracycline, 9-amino-8-chloro-5-hydroxy-6-

deoxy-4-dedimethylaminotetracycline, 9-(N,N-dimethyl) glycyllamino-5-hydroxy-6-deoxy-4-dedimethylaminotetracycline, 9-nitro-7-dimethylamino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 9-acetamido-7-dimethylamino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 9-(N, N-dimethyl) glycyllamino-7-dimethylamino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, and 9-ethoxythiocarbonylthio-7-dimethylamino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline.

In addition, the D ring may be halogenated at the C8 position to provide 8-halo-dedimethylaminotetracycline derivatives. As used in this specification, halogens can be chlorine, fluorine, bromine, and iodine. Some examples of 8-halo-dedimethylaminotetracycline derivatives are 9-amino-8-chloro-7-dimethylamino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 9-amino-8-chloro-7-dimethylamino-5-hydroxy-6-deoxy-4-dedimethylaminotetracycline and 9-amino-8-chloro-6-demethyl-6-deoxy-4-dedimethylaminotetracycline.

Another preferred method of the present invention provides for reducing the risk of cataract development in a mammal by administering a tetracycline derivative selected from the group consisting of 6a-benzylthiomethylenetetracycline, tetracyclinotrile, the mono-N-alkylated amide of tetracycline, 6-fluoro-6-demethyltetracycline, 11a-chlorotetracycline, tetracycline pyrazole and 12a-deoxytetracycline and its derivatives.

Another preferred method of the present invention provides that the tetracycline derivative is an antimicrobial tetracycline. Antimicrobial tetracycline derivatives which can be used with the methods of the present invention include tetracycline, minocycline and doxycycline.

Another method of the present invention provides for reducing the risk of cataract development in a mammal by administering a tetracycline derivative systemically. Preferably, the tetracycline derivative is administered systemically by a controlled release delivery system.

Additional methods of the present invention provide for reducing the risk of cataract development in a mammal by administering a tetracycline derivative orally or topically.

5 These and other advantages of the present invention will be appreciated from the detailed description and drawings which are set forth herein. The detailed description and the drawings enhance the understanding of the invention, but are not intended to limit the scope of the invention.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

10 Preferred embodiments of the invention have been chosen for purposes of illustration and description, but are not intended in any way to restrict the scope of the invention. The preferred embodiments of certain aspects of the invention are shown in the accompanying drawings, wherein:

15 Figure 1 is a bar graph illustration of the development of cataracts in untreated Type II ZDF/GMI diabetic rats and diabetic rats treated with doxycycline, CMT-3 (6-demethyl-6-deoxy-4-dedimethylaminotetracycline) and CMT-8 (5-hydroxy-6- $\alpha$ -deoxy-4-dedimethylaminotetracycline).

Figure 2 is a photographic illustration of the development of cataracts in untreated Type II ZDF/GMI diabetic rats in comparison with healthy control rats.

20 Figure 3 is a photographic illustration of the development of cataracts in Type II ZDF/GMI diabetic rats treated with doxycycline and CMT-8.

### **DETAILED DESCRIPTION OF THE INVENTION**


The present invention provides for methods of reducing the risk of cataract development in a mammal. The methods provide for administering to a mammal an

effective amount of a tetracycline derivative.


The methods of the present invention can be used in any living mammal. Mammals include, for example, humans, as well as pet animals such as dogs and cats, laboratory animals, such as rats and mice, and farm animals, such as horses and cows.

5

A tetracycline derivative as used herein, exhibits the following general ring structure:



The numbering system of the multiple ring nucleus is as follows:



10

Tetracycline, as well as the 5-OH (oxytetracycline, e.g. Terramycin) and 7-Cl (chlorotetracycline, e.g., Aureomycin) derivatives, exist in nature, and are all well known antibiotics. Semisynthetic tetracyclines include, for example, doxycycline, minocycline and methacycline.

15

A class of compounds has been defined which are structurally related to the antibiotic tetracyclines, but which have had their antibiotic activity substantially or completely eliminated by chemical modification. Substantial elimination of antibiotic activity occurs when the antibiotic activity is substantially less than that of tetracycline. Preferably, the antibiotic activity is at least approximately two times less than that of tetracycline, more preferably at least approximately five times less than

that of tetracycline, and even more preferably at least approximately ten times less than that of tetracycline.

The modifications that can and cannot be made to the basic tetracycline structure were reviewed by Mitscher, L.A., The Chemistry of the Tetracycline Antibiotics, Marcel Dekker, New York (1978), Ch. 6. According to Mitscher, the modification at positions 5-9 of the tetracycline ring system can be made without causing the complete loss of antibiotic properties. However, changes to the basic structure of the ring system, or replacement of substituents at positions 1-4 or 10-12, generally lead to synthetic tetracyclines with substantially less or essentially no antibacterial activity.

Chemically modified tetracyclines (CMT's) derivatives include, for example, 4-dedimethylaminotetracycline (CMT-1), tetracyclinonitrile (CMT-2), 6-demethyl-6-deoxy-4-dedimethylaminotetracycline (CMT-3), 7-chloro-4-dedimethylaminotetracycline (CMT-4), tetracyclinopyrazole (CMT-5), 4-hydroxy-4-dedimethylaminotetracycline (CMT-6), 12  $\alpha$ -deoxy-4-dedimethylaminotetracycline (CMT-7), 5-hydroxy-6- $\alpha$ -deoxy-4-dedimethylaminotetracycline (CMT-8), 4-dedimethylamino-12- $\alpha$ -deoxyanhydrotetracycline (CMT-9) and 7-dimethylamino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline (CMT-10). All are useful as non-antibacterial tetracyclines that reduce the risk of cataract development in a mammal.

A particularly preferred tetracycline derivative suitable for use according to the present invention is CMT-8.

Tetracycline derivatives which possess antibacterial activity are also contemplated by the present invention. However, such compounds are preferably employed in an amount which has substantially no antibacterial activity but which is effective for reducing the risk of cataract formation in a mammal. Preferred compounds of this type include tetracycline, minocycline and doxycycline.

The chemically modified and antimicrobial tetracycline derivatives can be made by methods known in the art. See, for example, Mitscher, L.A., *The Chemistry of the Tetracycline Antibiotics*, Marcel Dekker, New York (1978), Ch. 6 and U.S. Pat. Nos. 4,704,383 and 5,532,227.

5           According to the methods of the present invention, an effective amount of a tetracycline derivative is administered. An "effective amount" as used herein is that amount effective to achieve the specified result of reducing the risk of cataract formation. Preferably, the tetracycline derivative is provided in an amount which has little or no antimicrobial activity. A tetracycline derivative is not effectively  
10           antimicrobial if it does not significantly prevent the growth of microbes. Accordingly, the methods of the present invention can beneficially employ a tetracycline derivative which has been modified chemically to reduce or eliminate its antimicrobial properties. The use of CMT's is preferred in the present invention since they can be used at higher levels than antimicrobial tetracyclines, while avoiding  
15           certain disadvantages such as the indiscriminate killing of beneficial microbes and the emergence of resistant microbes, which often accompanies the use of antimicrobial or antibacterial amounts of such compounds.

          Tetracycline derivatives which are useful according to the methods of the present invention appear to exhibit their beneficial effect in a dose-dependent manner  
20           although CMT-8 exhibits its best effect when absorbed. Thus, within broad limits, administration of larger quantities of a tetracycline derivative has been observed to further reduce the risk of cataract formation than does administration of smaller amounts. Moreover, efficacy has been observed at dosages below the level at which toxicity is seen.

25           Maximal dosage for a subject is the highest dosage which does not cause undesirable or intolerable side effects. For example, the tetracycline compound can be administered in an amount of from about 0.1 mg/kg/day to about 30 mg/kg/day, and preferably from about 1 mg/kg/day to about 18 mg/kg/day. For the purpose of the

present invention, side effects can include clinically significant antimicrobial or antibacterial activity, as well as toxic effects. For example, a dose in excess of about 50 mg/kg/day would likely produce side effects in most mammals, including humans. In any event, the practitioner is guided by skill and knowledge in the field, and the present invention includes without limitation dosages which are effective to achieve the described effect.

The methods of the present invention involve administering a tetracycline derivative in an amount which is effective for reducing the risk of cataract formation in a mammal.

Administering tetracycline derivatives can be accomplished in a variety of ways. For example, tetracycline derivatives can be administered systemically by the parenteral and enteral routes, and include controlled release delivery systems. For example, a tetracycline derivative can easily be administered intravenously (e.g., intravenous injection) which is a preferred route of delivery. Intravenous administration can be accomplished by mixing the tetracycline derivative in a suitable pharmaceutical carrier (vehicle) or excipient as understood by practitioners in the art.

Oral or enteral use is also contemplated, and formulations such as tablets, capsules, pills, troches, elixirs, suspensions, syrups, wafers, chewing gum and the like can be employed to provide a tetracycline derivative.

Alternatively, delivery of a tetracycline derivative can include topical application. Accordingly, the carrier is preferably suited for topical use. Compositions deemed to be suited for such topical use include gels, salves, lotions, creams, ointments and the like. The tetracycline derivative can also be incorporated with a support base or matrix or the like to provide a pre-packaged surgical or burn dressing or bandage which can be directly applied to skin. Topical application of tetracycline derivatives in amounts of up to about 25% (w/w) in a vehicle are therefore appropriate. More preferably, application of tetracycline derivatives in

amounts of from about 0.1% to about 10% is believed to effectively reduce the risk of cataract formation according to the present invention. It is believed that these quantities do not induce significant toxicity in the subject being treated.

5 Combined or coordinated topical and systemic administration of tetracycline derivatives is also contemplated under the invention. For example, a non-absorbable tetracycline compound, such as CMT-2 or CMT-6, can be administered topically, while a tetracycline compound capable of substantial absorption and effective systemic distribution in a subject, such as CMT-1, CMT-3, CMT-7 or CMT-8, can be administered systemically.

10

The invention has been developed based on the unexpected observation by Applicants that tetracycline derivatives reduce the risk of cataract formation in a mammal. Applicants are unaware of any physiological or biochemical basis for expecting that tetracyclines would reduce the risk of cataract formation in a mammal. It is, therefore, surprising that tetracycline derivatives are found to reduce the risk of cataract formation in a mammal.

15

Diabetics develop posterior subcapsular cataracts, characterized by a central opacity in the nucleus and cortex. In studies on ZDF/Gmi-*fa* rats, for example, non-diabetic controls did not develop clinically-detectable cataracts, whereas 65% of the untreated Type II diabetic rats developed these ocular lesions by 5 months of age, an increase in incidence that was statistically significant ( $p < 0.001$ ). Of extreme interest, treatment with CMT-8 appeared to prevent the development of these cataracts at five months while the antimicrobial parent compound doxycycline was not as effective. In the current study, CMT-8 administration to the Type II diabetic rats reduced the incidence of cataract development by 37% at five months. In fact, this therapeutic effect may have been underestimated since it was an unexpected finding and, therefore, was not documented at the earlier time points. The mechanisms by which Type II (and Type I) diabetes causes the development of these cataracts is not yet understood.

20

25

Systemic administration of a tetracycline derivative such as CMT-8 should prevent the development of cataracts, which if left untreated can lead to blindness. The use of this therapeutic in individuals prone to cataract formation prophylactically could reduce the need for surgical intervention. In the case of surgical treatment, the use of CMT-8 could prevent recurrence of cataracts.

Figure 1 shows the development of cataracts in untreated Type II ZDF/GMI diabetic rats and Type II ZDF/GMI diabetic rats treated with different tetracycline analogues. Specifically, approximately 60% of untreated Type II ZDF/GMI diabetic rats developed cataracts whereas approximately 42% of Type II ZDF/GMI diabetic rats treated with CMT-8 developed cataracts.

Type II ZDF/GMI diabetic rats treated with other tetracycline analogues did not exhibit as low a percentage of cataract development in comparison to Type II ZDF/GMI diabetic rats treated with CMT-8. In particular, approximately 60% of Type II ZDF/GMI diabetic rats treated with doxycycline developed cataracts and approximately 70% of Type II ZDF/GMI diabetic rats treated with CMT-3 developed cataracts.

It is believed that the lack of efficacy for doxycycline and CMT-3 was due to the serum concentration achieved. Doxycycline, CMT-3 and CMT-8 were each administered at an oral dose of 15 mg/kg with rat pK studies revealing that equivalent oral doses result in very different peak serum concentrations. The peak serum concentration for doxycycline was determined to be 0.9 ug/ml and the peak serum concentration for CMT-3 was determined to be 4.6 ug/ml. The peak serum concentration, though, for CMT-8 was determined to be 15 ug/ml. Liu et al., The Lipophilicity and Pharmacokinetics of Different Chemically-Modified Tetracyclines (CMTs) In Vivo, Current Medicinal Chemistry, submitted for publication in September 1999. Clearly, administration of higher oral doses of CMT-3 and doxycycline would be needed to achieve serum concentrations similar to CMT-8.

Figures 2 and 3 provide a comparison of healthy untreated rats with untreated Type II ZDF/GMI diabetic rats and Type II ZDF/GMI diabetic rats treated with doxycycline and CMT-8. Cataracts do not appear to have developed in the CMT-8 treated diabetic rats as they appear very similar to the non-diabetic control rats.

- 5 Treating diabetic rats with doxycycline to prevent cataract development does not appear to be effective although it is believed that the lack of efficacy was due to the low serum concentration as discussed above.

- 10 Thus, while there have been described what are presently believed to be the preferred embodiments of the present invention, those skilled in the art will realize that other and further embodiments can be made without departing from the spirit of the invention, and it is intended to include all such further modifications and changes as come within the true scope of the claims set forth herein.

**WHAT IS CLAIMED IS:**

1. A method of reducing the risk of cataract development in a mammal comprising administering to the mammal an effective amount of a tetracycline derivative.

2. A method according to Claim 1, wherein said tetracycline derivative is a non-antimicrobial tetracycline.

3. A method according to Claim 1, wherein said tetracycline derivative is a dedimethylaminotetracycline.

4. A method according to Claim 3, wherein said dedimethylaminotetracycline is selected from the group consisting of 4-dedimethylaminotetracycline, 4-dedimethylamino-5-oxytetracycline, 4-dedimethylamino-7-chlorotetracycline, 4-hydroxy-4-dedimethylaminotetracycline, 5a,6-anhydro-4-hydroxy-4-dedimethylaminotetracycline, 6 $\alpha$ -deoxy-5-hydroxy-4-dedimethylaminotetracycline, 6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 4-dedimethylamino-12a-deoxytetracycline, 12 $\alpha$ -deoxy-4-deoxy-4-dedimethylaminotetracycline, 12a, 4 $\alpha$ -anhydro-4-dedimethylaminotetracycline, 7-dimethylamino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 5-hydroxy-6 $\alpha$ -deoxy-4-dedimethylaminotetracycline, 4-dedimethylamino-12 $\alpha$ -deoxyanhydrotetracycline and 4-dedimethylamino-11-hydroxy-12a-deoxytetracycline.

5. A method according to Claim 1, wherein said tetracycline derivative is 6 $\alpha$ -deoxy 5-hydroxy 4-dedimethylamino tetracycline.

6. A method according to Claim 1, wherein said tetracycline derivative is selected from the group consisting of 6a-benzylthiomethylenetetracycline, tetracyclino-trile, the mono-N-alkylated amide of tetracycline, 6-fluoro-6-demethyltetracycline, 11a-chlorotetracycline, tetracycline pyrazole, and 12a-deoxytetracycline and its derivatives.

7. A method according to Claim 1, wherein said tetracycline derivative is an antimicrobial tetracycline.
8. A method according to Claim 7, wherein said tetracycline derivative is tetracycline.
9. A method according to Claim 7, wherein said tetracycline derivative is minocycline.
10. A method according to Claim 7, wherein said tetracycline derivative is doxycycline
11. A method according to Claim 1, wherein said tetracycline derivative is administered systemically.
12. A method according to Claim 11, wherein said tetracycline derivative is administered systemically by a controlled release delivery system.
13. A method according to Claim 1, wherein said tetracycline derivative is administered orally.
14. A method according to Claim 1, wherein said tetracycline derivative is administered topically.
15. A method according to Claim 1, wherein said tetracycline derivative is a tetracycline compound of the formulae:

Structure A	or	Structure B
Structure C	or	Structure D

R7 is selected from the group consisting of hydrogen, amino, nitro, mono(lower alkyl) amino, halogen, di(lower alkyl)amino, ethoxythiocarbonylthio, azido, acylamino, diazonium, cyano, and hydroxyl;

R6-a is selected from the group consisting of hydrogen and methyl;

10 R6 and R5 are selected from the group consisting of hydrogen and hydroxyl;

R8 is selected from the group consisting of hydrogen and halogen;

R9 is selected from the group consisting of hydrogen, amino, azido, nitro, acylamino, hydroxy, ethoxythiocarbonylthio, mono(lower alkyl) amino, halogen, diazonium, di(lower alkyl)amino and  $\text{RCH}(\text{NH}_2)\text{CO}$ ;

15 R is hydrogen or lower alkyl; and pharmaceutically acceptable and unacceptable salts thereof; with the following provisos:

when either R7 and R9 are hydrogen then R8 must be halogen; and

when R6-a, R6, R5 and R9 are all hydrogen and R7 is hydrogen, amino, nitro, halogen, dimethylamino or diethylamino, then R8 must be halogen; and

20 when R6-a is methyl, R6 and R9 are both hydrogen, R5 is hydroxyl, and R7 is hydrogen, amino, nitro, halogen or diethylamino, then R8 is halogen; and

when R6-a is methyl, R6 is hydroxyl, R5, R7 and R9 are all hydrogen, then R8 must be halogen; and

25 when R6-a, R6 and R5 are all hydrogen, R9 is methylamino and R7 is dimethylamino, then R8 must be halogen; and

when R6-a is methyl, R6 is hydrogen, R5 is hydroxyl, R9 is methylamino and R7 is dimethylamino, then R8 must be halogen; and

when R6-a is methyl, R6, R5 and R9 are all hydrogen and R7 is cyano, then R8 must be halogen.

16. A method according to Claim 1, wherein said tetracycline derivative is a tetracycline compound of the formulae:

Structure E or Structure F or

Structure G or Structure H

5 wherein:

R7 is selected from the group consisting of hydrogen, amino, nitro, mono(lower alkyl) amino, halogen, and di(lower alkyl)amino, ethoxythiocarbonylthio, azido, acylamino, diazonium, cyano, and hydroxyl;

R6-a is selected from the group consisting of hydrogen and methyl;

10 R6 and R5 are selected from the group consisting of hydrogen and hydroxyl;

R4 is selected from the group consisting of NOH, N-NH-A, and NH-A, where A is a lower alkyl group;

R8 is selected from the group consisting of hydrogen and halogen;

15 R9 is selected from the group consisting of hydrogen, amino, azido, nitro, acylamino, hydroxy, ethoxythiocarbonylthio, mono(lower alkyl) amino, halogen, di(lower alkyl)amino and  $RCH(NH_2)CO$ ;

R is hydrogen or lower alkyl; and pharmaceutically acceptable and unacceptable salts thereof; with the following provisos:

20 when R4 is NOH, N-NH-alkyl or NH-alkyl and R7, R6-a, R6, R5, and R9 are all hydrogen, then R8 must be halogen; and

when R4 is NOH, R6-a is methyl, R6 is hydrogen or hydroxyl, R7 is halogen, R5 and R9 are both hydrogen, then R8 must be halogen; and

when R4 is N-NH-alkyl, R6-a is methyl, R6 is hydroxyl and R7, R5, R9 are all hydrogen, then R8 must be halogen; and

25 when R4 is NH-alkyl, R6-a, R6, R5 and R9 are all hydrogen, R7 is hydrogen, amino, mono(lower alkyl)amino, halogen, di(lower alkyl)amino or hydroxyl, then R8 must be halogen; and

30 when R4 is NH-alkyl, R6-a is methyl, R6 and R9 are both hydrogen, R5 is hydroxyl, and R7 is mono(lower alkyl)amino or di(lower alkyl)amino, then R8 must be halogen; and

when R4 is NH-alkyl, R6-a is methyl, R6 is hydroxy or hydrogen and R7, R5, and R9 are all be hydrogen, then R8 must be halogen.

17. A method according to Claim 1, wherein said tetracycline derivative is a 4-dedimethylaminotetracycline compound having general formulae (I) through (IV):

**General Formula (I)**

**Structure I**

5 wherein: R7, R8, and R9 taken together in each case, have the following meanings:

	R7	R8	R9
	azido	hydrogen	hydrogen
	dimethylamino	hydrogen	azido
	hydrogen	hydrogen	amino
10	hydrogen	hydrogen	azido
	hydrogen	hydrogen	nitro
	dimethylamino	hydrogen	amino
	acylamino	hydrogen	hydrogen
	hydrogen	hydrogen	acylamino
15	amino	hydrogen	nitro
	hydrogen	hydrogen	(N,N-dimethyl)glycylamino
	amino	hydrogen	amino
	hydrogen	hydrogen	ethoxythiocarbonylthio
	dimethylamino	hydrogen	acylamino
20	dimethylamino	hydrogen	diazonium
	dimethylamino	chloro	amino
	hydrogen	chloro	amino
	amino	chloro	amino
	acylamino	chloro	acylamino
25	amino	chloro	hydrogen
	acylamino	chloro	hydrogen
	mono alkylamino	chloro	amino
	nitro	chloro	amino
	dimethylamino	chloro	acylamino
30	dimethylamino	chloro	dimethylamino

and

**General Formula (II)**

**Structure J**

or

**Structure K or**

**Structure L**

or

**Structure M**

35 wherein: R7, R8, and R9 taken together in each case, have the following meanings:

	R7	R8	R9
	azido	hydrogen	hydrogen
	dimethylamino	hydrogen	azido
	hydrogen	hydrogen	amino
40	hydrogen	hydrogen	azido
	hydrogen	hydrogen	nitro
	dimethylamino	hydrogen	amino
	acylamino	hydrogen	hydrogen
	hydrogen	hydrogen	acylamino
45	amino	hydrogen	nitro
	hydrogen	hydrogen	(N,N-dimethyl)glycylamino
	amino	hydrogen	amino
	hydrogen	hydrogen	ethoxythiocarbonylthio
	dimethylamino	hydrogen	acylamino
50	hydrogen	hydrogen	diazonium
	hydrogen	hydrogen	dimethylamino
	diazonium	hydrogen	hydrogen
	ethoxythiocarbonylthio	hydrogen	hydrogen
	dimethylamino	chloro	amino
55	amino	chloro	amino
	acylamino	chloro	acylamino
	hydrogen	chloro	amino
	amino	chloro	hydrogen
	acylamino	chloro	hydrogen
60	mono alkyl amino	chloro	amino
	nitro	chloro	amino

and

### General Formula (III)

#### Structure N

65 wherein: R8 is hydrogen or halogen and R9 is selected from the group consisting of nitro, (N,N-dimethyl)glycylamino, and ethoxythiocarbonylthio; and

### General Formula (IV)

Structure O

or

Structure P

70 wherein: R7, R8, and R9 taken together in each case, have the following meanings:

	R7	R8	R9
	amino	hydrogen	hydrogen
	nitro	hydrogen	hydrogen
	azido	hydrogen	hydrogen
75	dimethylamino	hydrogen	azido
	hydrogen	hydrogen	amino
	hydrogen	hydrogen	azido
	hydrogen	hydrogen	nitro
	bromo	hydrogen	hydrogen
80	dimethylamino	hydrogen	amino
	acylamino	hydrogen	hydrogen
	hydrogen	hydrogen	acylamino
	amino	hydrogen	nitro
	hydrogen	hydrogen	(N,N-dimethyl)glycylamino
85	amino	hydrogen	amino
	diethylamino	hydrogen	hydrogen
	hydrogen	hydrogen	ethoxythiocarbonylthio
	dimethylamino	hydrogen	methylamino
	dimethylamino	hydrogen	acylamino
90	dimethylamino	chloro	amino
	amino	chloro	amino
	acylamino	chloro	acylamino
	hydrogen	chloro	amino
	amino	chloro	hydrogen
95	acylamino	chloro	hydrogen
	mono alkyl amino	chloro	amino
	nitro	chloro	amino

and pharmaceutically acceptable and unacceptable salts thereof.

18. A method of treating cataract formation in a mammal comprising administering to the mammal an effective amount of a tetracycline derivative.

19. A method according to Claim 18, wherein said tetracycline derivative is a non-antimicrobial tetracycline.

20. A method according to Claim 18, wherein said tetracycline derivative is a dedimethylaminotetracycline.

21. A method according to Claim 20, wherein said dedimethylaminotetracycline is selected from the group consisting of 4-dedimethylaminotetracycline, 4-dedimethylamino-5-oxytetracycline, 4-dedimethylamino-7-chlorotetracycline, 4-hydroxy-4-dedimethylaminotetracycline, 5a,6-anhydro-4-hydroxy-4-dedimethylaminotetracycline, 6  $\alpha$  -deoxy-5-hydroxy-4-dedimethylaminotetracycline, 6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 4-dedimethylamino-12a-deoxytetracycline, 12 $\alpha$  -deoxy-4-deoxy-4-dedimethylaminotetracycline, 12a, 4  $\alpha$  -anhydro-4-dedimethylaminotetracycline, 7-dimethylamino-6-demethyl-6-deoxy-4-dedimethylaminotetracycline, 5-hydroxy-6-  $\alpha$  -deoxy-4-dedimethylaminotetracycline, 4-dedimethylamino-12  $\alpha$  -deoxyanhydrotetracycline and 4-dedimethylamino-11-hydroxy-12a-deoxytetracycline.

22. A method according to Claim 18, wherein said tetracycline derivative is 6  $\alpha$  -deoxy 5-hydroxy 4-dedimethylamino tetracycline.

23. A method according to Claim 18, wherein said tetracycline derivative is selected from the group consisting of 6a-benzylthiomethylenetetracycline, tetracyclinotriole, the mono-N-alkylated amide of tetracycline, 6-fluoro-6-demethyltetracycline, 11a-chlorotetracycline, tetracycline pyrazole, and 12a-deoxytetracycline and its derivatives.

24. A method according to Claim 18, wherein said tetracycline derivative is an antimicrobial tetracycline.

25. A method according to Claim 24, wherein said tetracycline derivative is tetracycline.

26. A method according to Claim 24, wherein said tetracycline derivative is minocycline.

27. A method according to Claim 24, wherein said tetracycline derivative is doxycycline.


28. A method according to Claim 18, wherein said tetracycline derivative is administered systemically.

29. A method according to Claim 28, wherein said tetracycline derivative is administered systemically by a controlled release delivery system.

30. A method according to Claim 18, wherein said tetracycline derivative is administered orally.

31. A method according to Claim 18, wherein said tetracycline derivative is administered topically.

32. A method according to Claim 18, wherein said tetracycline derivative is a tetracycline compound of the formulae:



5           wherein:

R7 is selected from the group consisting of hydrogen, amino, nitro, mono(lower alkyl) amino, halogen, di(lower alkyl)amino, ethoxythiocarbonylthio, azido, acylamino, diazonium, cyano, and hydroxyl;

R6-a is selected from the group consisting of hydrogen and methyl;

10           R6 and R5 are selected from the group consisting of hydrogen and hydroxyl;

R8 is selected from the group consisting of hydrogen and halogen;

R9 is selected from the group consisting of hydrogen, amino, azido, nitro, acylamino, hydroxy, ethoxythiocarbonylthio, mono(lower alkyl) amino, halogen, diazonium, di(lower alkyl)amino and  $RCH(NH_2)CO$ ;

15           R is hydrogen or lower alkyl; and pharmaceutically acceptable and unacceptable salts thereof; with the following provisos:

when either R7 and R9 are hydrogen then R8 must be halogen; and

when R6-a, R6, R5 and R9 are all hydrogen and R7 is hydrogen, amino, nitro,

halogen, dimethylamino or diethylamino, then R8 must be halogen; and

when R6-a is methyl, R6 and R9 are both hydrogen, R5 is hydroxyl, and R7 is hydrogen, amino, nitro, halogen or diethylamino, then R8 is halogen; and

5 when R6-a is methyl, R6 is hydroxyl, R5, R7 and R9 are all hydrogen, then R8 must be halogen; and

when R6-a, R6 and R5 are all hydrogen, R9 is methylamino and R7 is dimethylamino, then R8 must be halogen; and

when R6-a is methyl, R6 is hydrogen, R5 is hydroxyl, R9 is methylamino and R7 is dimethylamino, then R8 must be halogen; and

10 when R6-a is methyl, R6, R5 and R9 are all hydrogen and R7 is cyano, then R8 must be halogen.

33. A method according to Claim 18, wherein said tetracycline derivative is a tetracycline compound of the formulae:

<b>Structure E</b>	<b>or</b>	<b>Structure F or</b>
<b>Structure G</b>	<b>or</b>	<b>Structure H</b>

5 wherein:

R7 is selected from the group consisting of hydrogen, amino, nitro, mono(lower alkyl) amino, halogen, and di(lower alkyl)amino, ethoxythiocarbonylthio, azido, acylamino, diazonium, cyano, and hydroxyl;

R6-a is selected from the group consisting of hydrogen and methyl;

10 R6 and R5 are selected from the group consisting of hydrogen and hydroxyl;

R4 is selected from the group consisting of NOH, N-NH-A, and NH-A, where A is a lower alkyl group;

R8 is selected from the group consisting of hydrogen and halogen;

15 R9 is selected from the group consisting of hydrogen, amino, azido, nitro, acylamino, hydroxy, ethoxythiocarbonylthio, mono(lower alkyl) amino, halogen, di(lower alkyl)amino and RCH(NH<sub>2</sub>)CO;

R is hydrogen or lower alkyl; and pharmaceutically acceptable and unacceptable salts thereof; with the following provisos:

- when R4 is NOH, N-NH-alkyl or NH-alkyl and R7, R6-a, R6, R5, and R9 are  
 20 all hydrogen, then R8 must be halogen; and
- when R4 is NOH, R6-a is methyl, R6 is hydrogen or hydroxyl, R7 is halogen,  
 R5 and R9 are both hydrogen, then R8 must be halogen; and
- when R4 is N-NH-alkyl, R6-a is methyl, R6 is hydroxyl and R7, R5, R9 are all  
 hydrogen, then R8 must be halogen; and
- 25 when R4 is NH-alkyl, R6-a, R6, R5 and R9 are all hydrogen, R7 is hydrogen,  
 amino, mono(lower alkyl)amino, halogen, di(lower alkyl)amino or hydroxyl, then R8  
 must be halogen; and
- when R4 is NH-alkyl, R6-a is methyl, R6 and R9 are both hydrogen, R5 is  
 hydroxyl, and R7 is mono(lower alkyl)amino or di(lower alkyl)amino, then R8 must  
 30 be halogen; and
- when R4 is NH-alkyl, R6-a is methyl, R6 is hydroxy or hydrogen and R7, R5,  
 and R9 are all be hydrogen, then R8 must be halogen.

34. A method according to Claim 18, wherein said tetracycline derivative  
 is a 4-dedimethylaminotetracycline compound having general formulae (I) through  
 (IV):

#### General Formula (I)

5

#### Structure I

wherein: R7, R8, and R9 taken together in each case, have the following meanings:

	R7	R8	R9
	azido	hydrogen	hydrogen
	dimethylamino	hydrogen	azido
10	hydrogen	hydrogen	amino
	hydrogen	hydrogen	azido
	hydrogen	hydrogen	nitro
	dimethylamino	hydrogen	amino
	acylamino	hydrogen	hydrogen
15	hydrogen	hydrogen	acylamino
	amino	hydrogen	nitro
	hydrogen	hydrogen	(N,N-dimethyl)glycylamino
	amino	hydrogen	amino
	hydrogen	hydrogen	ethoxythiocarbonylthio
20	dimethylamino	hydrogen	acylamino
	dimethylamino	hydrogen	diazonium

	dimethylamino	chloro	amino
	hydrogen	chloro	amino
	amino	chloro	amino
25	acylamino	chloro	acylamino
	amino	chloro	hydrogen
	acylamino	chloro	hydrogen
	mono alkylamino	chloro	amino
	nitro	chloro	amino
30	dimethylamino	chloro	acylamino
	dimethylamino	chloro	dimethylamino

and

General Formula (II)

	Structure J	or	Structure K or
35	Structure L	or	Structure M

wherein: R7, R8, and R9 taken together in each case, have the following meanings:

	R7	R8	R9
	azido	hydrogen	hydrogen
	dimethylamino	hydrogen	azido
40	hydrogen	hydrogen	amino
	hydrogen	hydrogen	azido
	hydrogen	hydrogen	nitro
	dimethylamino	hydrogen	amino
	acylamino	hydrogen	hydrogen
45	hydrogen	hydrogen	acylamino
	amino	hydrogen	nitro
	hydrogen	hydrogen	(N,N-dimethyl)glycylamino
	amino	hydrogen	amino
	hydrogen	hydrogen	ethoxythiocarbonylthio
50	dimethylamino	hydrogen	acylamino
	hydrogen	hydrogen	diazonium
	hydrogen	hydrogen	dimethylamino
	diazonium	hydrogen	hydrogen
	ethoxythiocarbonylthio	hydrogen	hydrogen
55	dimethylamino	chloro	amino
	amino	chloro	amino
	acylamino	chloro	acylamino
	hydrogen	chloro	amino
	amino	chloro	hydrogen
60	acylamino	chloro	hydrogen

	mono alkyl amino	chloro	amino
	nitro	chloro	amino
and			

**General Formula (III)**

65

**Structure N**

wherein: R8 is hydrogen or halogen and R9 is selected from the group consisting of nitro, (N,N-dimethyl)glycylamino, and ethoxythiocarbonylthio; and

**General Formula (IV)**

70

**Structure O**

or

**Structure P**

wherein: R7, R8, and R9 taken together in each case, have the following meanings:

	R7	R8	R9
	amino	hydrogen	hydrogen
	nitro	hydrogen	hydrogen
75	azido	hydrogen	hydrogen
	dimethylamino	hydrogen	azido
	hydrogen	hydrogen	amino
	hydrogen	hydrogen	azido
	hydrogen	hydrogen	nitro
80	bromo	hydrogen	hydrogen
	dimethylamino	hydrogen	amino
	acylamino	hydrogen	hydrogen
	hydrogen	hydrogen	acylamino
	amino	hydrogen	nitro
85	hydrogen	hydrogen	(N,N-dimethyl)glycylamino
	amino	hydrogen	amino
	diethylamino	hydrogen	hydrogen
	hydrogen	hydrogen	ethoxythiocarbonylthio
	dimethylamino	hydrogen	methylamino
90	dimethylamino	hydrogen	acylamino
	dimethylamino	chloro	amino
	amino	chloro	amino
	acylamino	chloro	acylamino
	hydrogen	chloro	amino
95	amino	chloro	hydrogen

acylamino  
mono alkyl amino  
nitro





chloro  
chloro  
chloro


hydrogen  
amino  
amino

100


and pharmaceutically acceptable and unacceptable salts thereof.

## STRUCTURES A-P







Structure E




Structure F




Structure G




Structure H




Structure I




Structure J




Structure K




Structure L




Structure M



Structure N



Structure O



Structure P

FIG. 1




FIG. 2





FIG. 3



CMT-8 TREATED DIABETIC

DOXYCYCLINE TREATED DIABETIC

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>A61K 31/65</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 00/18353</b> <b>(43) International Publication Date:</b> 6 April 2000 (06.04.00)
<b>(21) International Application Number:</b> PCT/US99/22354 <b>(22) International Filing Date:</b> 28 September 1999 (28.09.99) <b>(30) Priority Data:</b> 60/102,056 28 September 1998 (28.09.98) US <b>(71) Applicant (for all designated States except US):</b> THE RE-SEARCH FOUNDATION OF STATE UNIVERSITY OF NEW YORK [US/US]; P.O. Box 9, Albany, NY 12201-0009 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> RYAN, Maria, Emanuel [US/US]; 5 Vannina Place, Huntington, NY 11743 (US). GOLUB, Lorne, M. [US/US]; 29 Whitney Gate, Smith-town, NY 11787 (US). RAMAMURTHY, Nungavaram, S. [US/US]; 10 Lynam Court, Smithtown, NY 11787 (US). <b>(74) Agent:</b> BARON, Ronald, J.; Hoffmann & Baron, LLP, 6900 Jericho Turnpike, Syosset, NY 11791 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <b>(88) Date of publication of the international search report:</b> 6 July 2000 (06.07.00)
<b>(54) Title:</b> A NOVEL INHIBITOR OF CATARACT FORMATION		
<b>(57) Abstract</b>  Methods of reducing the risk of cataract development in a mammal are provided and include administering to the mammal an effective amount of a tetracycline derivative. A preferred tetracycline derivative administered according to the methods of the present invention is 6 $\alpha$ -deoxy 5-hydroxy-4-dedimethylaminotetracycline.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/22354

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/65  
US CL : 514/152, 912

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : NONE

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
CAS-ONLINE, USPATFULL

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,244,673 A (GEJKOVA et al.) 14 September 1993, see the entire document.	1-34
X	US 5,474,764 A (PATEL et al.) 12 December 1995, see the entire document.	1-34
X, P	US 5,814,655 A (PATEL et al.) 29 September 1998, see the entire document.	1-34
X, P	US 5,929,055 A (RYAN et al.) 27 July 1999, see the entire document.	1-34

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

17 MARCH 2000

Date of mailing of the international search report

04 APR 2000

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

KEVIN E. WEDDINGTON

Telephone No. (703) 308-1235